

AD \_\_\_\_\_

MIPR NO: 94MM4586

TITLE: CESSATION OF FOCAL EPILEPTIFORM ACTIVITY USING  
CORTICAL CALCIUM BLOCKER

SUBTITLE: Amelioration of Focal Epileptiform Activity Using  
a Topical (Cortical) Agent (Nimodipine) in a  
Porcine Model

PRINCIPAL INVESTIGATOR: William C. Bergman, COL, MC, USAR

CONTRACTING ORGANIZATION: Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, Maryland 20814-4799

REPORT DATE: January 31, 1995

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick  
Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19950222 006

DTIC  
ELECTE  
FEB 28 1995  
S G D

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 31 Jan 95	3. REPORT TYPE AND DATES COVERED Final: 08 Aug 94 - 31 Dec 94	
4. TITLE AND SUBTITLE Cessation of Focal Epileptiform Activity Using Cortical Calcium Blocker			5. FUNDING NUMBERS MIPR No. 94MM4586	
6. AUTHOR(S) William C. Bergman, COL, MC Leon E. Moores, MAJ, MC				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Uniformed Services University of the Health Sciences 4301 Jones Bridge Road Bethesda, MD, 20814-4799			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick Frederick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Subtitle: Amelioration of Focal Epileptiform Activity Using a Topical (Cortical) Agent (Nimodipine) in a Porcine Model				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) 60-80 pound male pigs were subjected to general anesthesia, using a balanced narcotic technique (fentanyl). The animals were then subjected to removal of the entire flat surface of the calvarium, as well as the dura (aside from the superior sagittal sinus). A cortical electroencephalographic (EEG) grid (PMT Corp.) was then placed over the surface of the exposed brain. Through a 3mm. hole in the grid, a seizure focus was established, using penicillin (5,000,000 U./cc.). Once this was definitively established, the penicillin was removed and replaced with a calcium channel blocker (nimodipine). In two out of four procedures, the focal epileptiform activity was ameliorated or altered.				
14. SUBJECT TERMS anticonvulsant      epileptiform discharge      nimodipine pig      topical			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

DTIC QUALITY INSPECTED 1

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

N.A. Where copyrighted material is quoted, permission has been obtained to use such material.

V.A. Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

w.B. Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

w.B. In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).


N.A. For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N.A. In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N.A. In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N.A. In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

 2a,mc 28 Jan '95  
PI - Signature Date

## INTRODUCTION

Although the vast majority of patients with seizure disorders can be adequately managed with systemic (oral) anticonvulsant agents, a small but significant minority remain intractable to such measures. Some of these patients eventually require ablative surgery (resection of small volumes of brain tissue) in order to eventually control their seizures. The extent of resection in such surgical procedures is based on a combination of preoperative and intraoperative EEG monitoring, but the exact dimensions and limits of the brain to be resected, in order to effect the desired result yet not injure any needed brain, is, at best, less than exact. It would be highly beneficial to have a technique which would allow reversible seizure control which could be performed preoperatively, or even as a separate surgical procedure, prior to removing brain and creating a situation which, if the wrong judgment regarding dimensions is made, leaves the patient with either an inadequate resection or irreversible loss of function. A topical anticonvulsant agent (a substance which could be applied directly to the cortex) which could be delivered using a device, the components of which are already in existence, would be highly beneficial in this regard.

An additional benefit of such an agent would be its potential use in patients with open or penetrating head injuries. Such patients, which are all too commonly seen in the combat scenario, will have exposed brain, and are very likely to seize. This is even more likely if the patient has been exposed to certain chemical agents. The "golden opportunity" to most effectively and definitively control these seizures is as early as possible, potentially at the scene of the trauma. Although systemic anticonvulsant agents are commonly given at this point in this scenario, they are frequently inadequate to effect the desired results. A topical agent which could be applied directly to the cortex, in such a situation, might well be highly beneficial.

For the reasons described above, we are endeavoring to develop an agent which will control seizure activity when applied directly to the brain cortex surface.

Work by Lockard (1) suggests that a calcium channel blocker flunarizine applied directly to the cortex of Rhesus monkeys, which had been subjected to the development of an epileptic focus (using alumina gel) showed a significant diminution of the frequency and intensity of their seizures. Previous work by this author (2) suggests that a similar calcium channel blocker (nimodipine) will ameliorate focal epileptiform discharges, when applied directly to the cortex of pigs subjected to a penicillin focus. This latter work proved to be somewhat sporadically repeatable, suggesting that an optimal technique needed further development. To this end, we are endeavoring to improve and perfect this technique.

## MATERIALS AND METHODS

60 to 80 lb. male (in order to avoid hormonal variations) pigs were subjected to general anesthesia, using a balanced narcotic (fentanyl) technique. The skin was incised in the mid-sagittal line along the entire A-P diameter of the calvarium. After hemostasis was achieved, the scalp was elevated away from the entire dorsal surface of the calvarium, and that bony structure was removed in its entirety. Again, hemostasis was achieved, and the exposed dura was then removed, bilaterally, except for the superior sagittal sinus. A 16-lead cortical EEG grid (PMT Corporation) was placed directly over the exposed brain, in such a manner that eight leads were present over the surface of cortex of each hemisphere. This was then connected to a standard electroencephalographic (EEG) recording device (Nicolet Corporation). Prior to its placement over the cortex, a 3mm hole was placed in the grid, through which various agents could be delivered.

After recording baseline tracings, a cotton pledget soaked in penicillin (5 million units per milliliter) was placed directly over the hole. The EEG tracing was observed for evidence of epileptiform electrical activity, and the pledget was removed as soon as the first spikes were noted. The cortex was irrigated, in order to remove all traces of penicillin, and the focus was allowed to develop. It was observed for approximately one-half hour, in order to ensure that the presence of the penicillin was no longer necessary.

A similar pledget, this time soaked in nimodipine (Nimotop R, Miles Pharmaceutical) was placed over the same hole. Results were then recorded.

At the end of the procedure, the animal was euthanatized by overdose.

## RESULTS

Five subject animals were subjected to the described procedures. One procedure was aborted because the animal had a preexistent generalized seizure discharge noted when the grid was first applied. Of the remaining four, two developed seizure foci which expanded well beyond the limits of the focus of the cortex under the hole. Of the remaining two, one developed burst suppression following the application of the nimodipine. This took place in spite of the fact that the focus did not appear to be deteriorating. In the remaining subject animal, the epileptiform focus was significantly diminished in its amplitude and frequency of discharge after the application of the nimodipine.

## CONCLUSIONS

1. The use of fentanyl, as opposed to the isofluorane used in the author's previous work (2), eliminated the presence of sporadic, preexistent spikes, which were a problem in the author's previous work.
2. Nimodipine ameliorates epileptiform activity generated by penicillin, but only if the epileptiform focus is limited to the area in direct contact with the nimodipine. This amelioration can be evidenced by either the diminution of epileptiform activity or the development of burst suppression.
3. Further work is necessary to show that these results can be repeated statistically.

## REFERENCES

1. Lockard, J.S.; Levy, R.H.; Congdon, W.C.; McNabb, M.L.: Flunarizine Administered Focally Delays Epileptogenesis in an Experimental Model. Epilepsia 30, 5, 1989, 649.
2. Bergman, W.C.; Baker, J.T.; Vance, S.C.; Barbaro, N.M.; Gelinas-Sorell, D.; Brooks, D.E.; Fredericks, D.; Rodkey, W.G.: Cessation of Focal Seizure Activity Using Calcium Channel Blocker (nimodipine) Applied Directly to the Cortex. Poster presented to the Annual Conference of the American Association of Neurological Surgeons, April 1991.